



## **Model of Ciprofloxacin Killing Enhanced by Hyperbaric Oxygen Treatment in *Pseudomonas aeruginosa* Biofilms**

**Gade, P. A.; Olsen, T. B.; Jensen, P. Ø.; Kolpen, Mette; Henneberg, Kaj-Åge; Sams, Thomas**

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Gade, P. A., Olsen, T. B., Jensen, P. Ø., Kolpen, M., Henneberg, K-Å., & Sams, T. (2018). *Model of Ciprofloxacin Killing Enhanced by Hyperbaric Oxygen Treatment in Pseudomonas aeruginosa Biofilms*. Poster session presented at 8th ASM Conference on Biofilms, Washington, District of Columbia, United States.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# 8th ASM Conference on Biofilms

October 7 – 11, 2018

Washington, DC



AMERICAN  
SOCIETY FOR  
MICROBIOLOGY

[Print this Page for Your Records](#)

[Close Window](#)

**Control/Tracking Number:** 2018-A-287-ASM

**Activity:** Abstract

**Current Date/Time:** 8/24/2018 4:01:40 AM

## Model of Ciprofloxacin Killing Enhanced by Hyperbaric Oxygen Treatment in *Pseudomonas aeruginosa* Biofilms

**Author Block:** P. A. Gade<sup>1</sup>, T. B. Olsen<sup>2</sup>, P. Ø. Jensen<sup>3</sup>, M. Kolpen<sup>3</sup>, N. Høiby<sup>3</sup>, K. Henneberg<sup>4</sup>, T. Sams<sup>4</sup>;

<sup>1</sup>Dept. of Applied Mathematics and Computer Science, Technical University of Denmark, Lyngby, DENMARK, <sup>2</sup>Dept. of Applied Mathematics and Computer Science, Lyngby, DENMARK, <sup>3</sup>Dept. of Clinical Microbiology, Rigshospitalet and Costerton Biofilm Center, Dept. of Immunology and Microbiology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, DENMARK, <sup>4</sup>Biomedical Engineering, Technical University of Denmark, Lyngby, DENMARK.

### Abstract:

In chronic *Pseudomonas aeruginosa* (PA) biofilm lung infections the bacteria are protected from the immune system of the host and from antibiotic treatment. It has been demonstrated that the susceptibility of the bacteria to antibiotic treatment can be enhanced by hyperbaric oxygen treatment. Here we present a reaction-diffusion model that describes the combined effect of ciprofloxacin diffusion, oxygen diffusion and depletion, bacterial growth and killing, and adaptation of the bacteria to ciprofloxacin. In the model, the oxygen diffusion and depletion use a set of parameters derived from experiments. The description of ciprofloxacin killing uses parameter values from the literature in combination with our estimates. The complete oxygen model comprises a reaction-diffusion equation describing the oxygen consumption by using a Michaelis-Menten reaction term. The oxygen model performed well in predicting oxygen concentrations in both time and depth into the biofilm. At 2.8 bar pure oxygen pressure, HBOT increases the penetration depth of oxygen into the biofilm by a factor of 4 and we see that hyperbaric oxygen treatment significantly increases the killing by ciprofloxacin in a PAO1 biofilm in alignment with the experimental results. References: PLoS ONE 13(6) 2018: e0198909. Antimicrobial Agents and Chemotherapy 61(9) 2017: AAC.01024-17. International Journal of Antimicrobial Agents 47(2) 2016: 163-167.

:

**Topic (Complete):** Biofilm: From Nature to Models

**Presentation Preference (Complete):** Poster or Oral presentation

**Additional Info (Complete):**

**Level of education?:** Ph.D.

**Travel Grants:** No

**Gender:** Male

**Minority:** No

**Status:** Complete

[American Society for Microbiology](#)

1752 N Street N.W.

Washington, D.C. 20036

Phone: (202) 737-3600

[Leave cOASIS Feedback](#)

Powered by [cOASIS](#), The Online Abstract Submission and Invitation System <sup>SM</sup>

© 1996 - 2018 [CTI Meeting Technology](#) All rights reserved. [Privacy Policy](#)

# Model of Ciprofloxacin Killing Enhanced by Hyperbaric Oxygen Treatment in *Pseudomonas aeruginosa*

Peter Alexander Vistar Gade<sup>1,2</sup>, Terkel Bo Olsen<sup>1,2</sup>,  
Peter Østrup Jensen<sup>3,4</sup>, Mette Kolpen<sup>3,4</sup>, Niels Højby<sup>3,4</sup>,  
Kaj-Åge Henneberg<sup>1</sup>, **Thomas Sams**<sup>1</sup>

1 Biomedical Engineering, Technical University of Denmark

2 Dept. of Applied Mathematics and Computer Science, Technical University of Denmark


3 Department of Clinical Microbiology, Rigshospitalet, Copenhagen, Denmark

4 Costerton Biofilm Center, Department of Immunology and Microbiology, University of Copenhagen

In chronic *Pseudomonas aeruginosa* (PA) biofilm lung infections the bacteria are protected from the immune system of the host and from antibiotic treatment. The effect of antibiotic treatment can be enhanced by hyperbaric oxygen treatment. We present a reaction-diffusion model that describes the combined effect of ciprofloxacin diffusion, oxygen diffusion and depletion, bacterial growth and killing, and adaptation of the bacteria to ciprofloxacin in a 2.8 bar oxygen treatment scheme.

PLoS ONE 13(6) 2018 e0198909  
Antimicrobial Agents and Chemotherapy 61(9) 2017: AAC.01024-17  
Int. J. Antimicrobial Agents 47(2) 2016: 163-167

## Survivor profile:



**Bacterial killing along the depth dimension of the biofilm.** 4 hour treatment scheme with six different doses of ciprofloxacin in a 5 mm biofilm model. The initial ciprofloxacin concentration in the supernatant is indicated, the equilibrated concentration is 5 times lower. [PLoS ONE 13(6) e0198909]

## Ciprofloxacin model with O<sub>2</sub> consumption:

Oxygen diffusion and consumption:

$$\frac{\partial c}{\partial t} = D_{O_2} \frac{\partial^2 c}{\partial z^2} - R_{\max} \frac{c}{K_m + c} \frac{\varepsilon}{\varepsilon_{\text{norm}}}$$

Ciprofloxacin diffusion:

$$\frac{\partial u}{\partial t} = D_{\text{cip}} \frac{\partial^2 u}{\partial z^2}$$

Bacterial growth and killing:

$$\frac{\partial \varepsilon}{\partial t} = \mu(c) \left(1 - \frac{\varepsilon}{\varepsilon_{\max}}\right) \varepsilon - K_{\max} \frac{\mu(c)}{\mu_{\max}} \frac{u^\gamma}{KC_{50S}^\gamma + u^\gamma} \varepsilon$$

Adaptation:

$$\frac{\partial \beta}{\partial t} = \left(\frac{S_{\max} u}{SC_{50} + u} - \beta\right) k_{\text{out}}$$

With:

$$KC_{50S} = KC_{50, \text{base}} (1 + \beta)$$

$$\mu(c) = \mu_{\max} \frac{c}{K_m + c}$$